

11-2017

## Predictors of In Hospital Mortality In Acute Pancreatitis

David E. Mossad

Wright State University, david.mossad@wright.edu

Brian V. Dinh

Wright State University, dinh.6@wright.edu

Ronald J. Markert

Wright State University - Main Campus, ronald.markert@wright.edu

Mustafa N. Musleh

Wright State University, mustafa.musleh@wright.edu

Sangeeta Agrawal

Wright State University, sangeeta.agrawal@wright.edu

Follow this and additional works at: [https://corescholar.libraries.wright.edu/internal\\_medicine](https://corescholar.libraries.wright.edu/internal_medicine)



Part of the [Internal Medicine Commons](#)

---

### Repository Citation

Mossad, D. E., Dinh, B. V., Markert, R. J., Musleh, M. N., & Agrawal, S. (2017). Predictors of In Hospital Mortality In Acute Pancreatitis. *Journal Pancreas*, 18 (6), 465-469.  
[https://corescholar.libraries.wright.edu/internal\\_medicine/192](https://corescholar.libraries.wright.edu/internal_medicine/192)

This Article is brought to you for free and open access by the Internal Medicine at CORE Scholar. It has been accepted for inclusion in Internal Medicine Faculty Publications by an authorized administrator of CORE Scholar. For more information, please contact [library-corescholar@wright.edu](mailto:library-corescholar@wright.edu).

# Predictors of In-Hospital Mortality in Acute Pancreatitis

David Mossad, Brian Dinh, Ronald Markert, Mustafa Musleh, Sangeeta Agrawal

Boonshoft School of Medicine, Wright State University, USA

## ABSTRACT

**Background and Aims** Pancreatitis presents with significant risk for morbidity and mortality acutely. The purpose of this study was to evaluate demographics, comorbidity burden, and hospital characteristics and their effect on in-hospital mortality in patients with acute pancreatitis. **Methods** Data was collected from The National Hospital Discharge Survey for patients admitted for acute pancreatitis from 2001-2010. Patients were separated into those who suffered in-hospital mortality and those that were discharged. Data was analyzed regarding demographics, length of stay, comorbidity burden, mortality, principal payment, and hospital size. Statistical comparisons were made using SPSS with chi-square and student T tests with a significance level of 0.05. **Results** 27,259 cases of acute pancreatitis were identified, with 573 who suffered in-hospital mortality. The mortality group was significantly older and had a longer length of stay. Patients with four or more comorbidities had a higher mortality rate (2.5%) compared to those with three or less (0.1%) ( $p < 0.001$ ). Region displayed a difference in mortality, with the Midwest showing a lower rate (1.6%) compared to all other regions. Mortality was lowest in small hospitals. Hospital ownership, however, did not show any significant differences in mortality. There was a difference in mortality rate when comparing principal forms of payment ( $p < 0.001$ ), with Medicare and Medicaid patients having the highest mortality rates. **Conclusions** Pancreatitis poses significant implications in morbidity and mortality in the acute setting. This study shows that factors such as age, length of stay, comorbidity burden, principal source of payment, and regional demographics have an impact on in-hospital mortality.

## INTRODUCTION

Acute pancreatitis (AP) is an inflammatory process of the pancreas that affects approximately 4.9 to 35 per 100,000 in the general population [1]. It is the leading cause of gastrointestinal hospital admission [2]. A large epidemiological study found that between 1988 and 2003, the mortality from acute pancreatitis decreased from 12% to 2% with a higher incidence of mortality among black males in their early 50's [3]. Overall mortality of patients with acute pancreatitis admitted to the hospital is estimated to be 5% to 10% varying from 3% in mild AP to 30% in severe AP [4, 5].

With significant risk of morbidity and mortality associated with acute pancreatitis, multiple scoring systems have been developed to assess its severity. However, no single scoring system has been shown to be superior to others in predicting in-hospital mortality [6, 7]. One of the early systems, Ranson's criteria, focuses on 11 factors with 5 assessed at the time of admission and 6 assessed 48 hours into admission. For scores  $< 3$ , mortality was estimated to be 0% to 3%, scores  $\geq 3$  were associated with a 11% to 15% mortality, and scores  $\geq 6$

were associated with a 40% mortality. Although used widely, a meta-analysis showed that Ranson's criteria had poor predictive value for acute pancreatitis severity [8].

Acute Physiology and Chronic Health Examination (APACHE) II, another cumbersome scoring system, was developed for use with critically ill patients, limiting its applicability. The APACHE II score uses age and 12 physiologic measures and can be calculated each day. Several studies have shown that an APACHE II score of less than 8 has a mortality rate  $< 4\%$  while a score greater than 8 results in mortality between 11% to 18% [5, 9].

Bedside Index of Severity in Acute Pancreatitis (BISAP) is a simpler calculation for determining early in-hospital mortality of patients admitted for acute pancreatitis. A BISAP score uses point assignments: 1 point for each of the following - BUN  $> 25$  mg/dL, abnormal mental status with Glasgow Coma Scale score  $< 15$ , evidence of Systemic Inflammatory Response Syndrome (SIRS),  $> 60$  years of age, and imaging revealing pleural effusion. Patients with a score of zero have less than 1% mortality while a score of 3 to 5 is associated with  $> 15\%$  mortality.

A recent validation study found that BISAP had similar performance to APACHE II and Ranson's criteria [10]. Moreover, BISAP compares favorably with newer scoring systems for bedside AP assessment, such as the "harmless acute pancreatitis score" and the "CT severity index" [10, 11].

In sum, AP severity scores are often unwieldy to measure and may require recalculation at later times during the hospital admission. In contrast, rather than focus on an

Received September 07th, 2017 - Accepted October 10th, 2017

**Keywords** Pancreatitis; Morbidity; Mortality

**Abbreviations** SPSS Statistical Package for the Social Sciences

**Correspondence** Sangeeta Agrawal

Division of Gastroenterology

Weber CHE Building, 128 E Apple St, 2nd Floor

Dayton, OH 45409

**Tel** +937-208-2124

**Fax** +937-208-2621

**E-mail** sagrawalgi@gmail.com

AP score or index, other studies have investigated patient demographics (e.g., age), clinical characteristics (e.g., BMI, SIRS), various laboratory values, and radiological findings to determine AP severity and risk of mortality [12, 13, 14].

We examined selected demographic factors, comorbidities, and health care system characteristics that might be predictors of mortality in hospitalized patients with acute pancreatitis.

## METHODS

Study data was acquired from the National Hospital Discharge Survey (NHDS), which is collected every year through the National Center for Health Statistics.

The NHDS collects patient hospitalizations and discharges from nonfederal hospitals across the United States with average length of stay of <30 days and  $\geq 6$  beds. Patient data in the NHDS is de-identified and made available to the public via the Center for Disease Control, making it exempt from an Institutional Review Board. The NHDS database was searched using International Classification of Diseases – Ninth Revision (ICD-9) diagnosis codes for patients admitted to hospitals with acute pancreatitis (ICD-9 code 577.0) from 2001-2010. For all patients included in the study, the following data was obtained: patient demographics (age, sex, race, marital status), hospital length of stay (LOS), hospital size (by number of beds), location/ownership (proprietary, government, or church/nonprofit), type of admission (emergency/urgent/elective), source of admission, comorbidity burden, and mortality. This cohort was separated into two groups: those who died during the hospitalization and those who survived the hospitalization.

Means and standard deviations are reported for continuous variables, and counts and percents for categorical variables. The independent samples Mann-Whitney Test was used for comparisons involving the two groups (mortality vs. no mortality) and a second variable measured on a continuous scale. The chi square test was used to compare the two groups on categorical variables. Inferences for these univariate statistical tests were made at the 0.05 level of significance with no corrections for multiple comparisons. Multivariable logistic regression was used to identify independent risk factors of mortality. Data was first recorded in Excel, and analyses were conducted using IBM SPSS Statistics 23.0 (IBM, Armonk, NY).

## RESULTS

**Table 1** shows the results for 27,259 patients admitted for acute pancreatitis during the 2001-2010 periods. Of this group, 573 (2.1%) suffered in-hospital mortality. The mean age of patients who died was 64.5 years, which was significantly higher than the survival group's mean age of 51.8 years ( $p < 0.001$ ). Only two races had a large number of patients: white ( $n = 13,610$ ) and African-American/black ( $n = 4984$ ), and there was no difference in mortality between these two groups (white 2.3% vs. A-A/black 2.2%;  $p = 0.55$ ).

Mean length of stay was nearly double (12.0 vs. 6.4 days,  $p < 0.001$ ) in the mortality group. Patients with four or more comorbidities had a significantly higher mortality rate (2.5%) compared to those with three or fewer comorbidities (0.1%) ( $p < 0.001$ ).

Mortality was greater as hospital size increased, with the largest hospitals having the highest mortality rate (2.6%) and the smallest hospitals having the lowest mortality rate (1.3%) ( $p < 0.001$ ). Mortality did not differ by type of hospital ownership ( $p = 0.49$ ). Of the 10 principal forms of payment, Medicare patients had significantly higher mortality (3.6%) than patients with other insurers – e.g., Medicaid (2.0%), HMO/PPO (1.4%), and self-pay (1.0%) ( $p < 0.001$ ).

Mortality did not differ by type of admission (emergency  $n = 18,382$ ; urgent  $n = 4,287$ ; and elective  $n = 1,729$ ). Elective patients had a mortality rate of 2.8%, emergency 2.1%, and urgent 1.8% ( $p = 0.06$ ). The hospitals in the Midwest had a lower mortality rate (1.6%) compared to the Northeast (2.5%), South (2.2%), and West (2.4%) ( $p = 0.002$ ).

Multivariable logistic regression (MLR) analysis was used to identify independent predictors of mortality. Predictors entered into the MLR equation were: age, sex, race (white, African-American, other), comorbidities (3 or fewer, 4 or more), admission type (emergency, urgent, elective), hospital length of stay, hospital region (Midwest, other 3 regions), hospital size (99 or fewer beds, 100 or more beds), hospital ownership (proprietary, government, church/nonprofit), and principal payment (Medicare, 9 other insurers)

**Table 2** shows the six independent risk factors. The highest adjusted odds ratio (AOR) was four or more comorbidities (AOR=15.22 [95% CI=4.87-47.54];  $p < 0.001$ ), followed by hospital size of 100 beds or more (AOR=1.42 [95% CI=1.04-1.95];  $p = 0.029$ ), Midwest region (AOR=1.31 [95% CI=1.05-1.64];  $p = 0.017$ ), male sex (AOR=1.26 [95% CI=1.05-1.52];  $p = 0.015$ ), increasing age (AOR=1.03 [95% CI=1.03-1.04];  $p < 0.001$ ), and longer hospital LOS (AOR=1.03 [95% CI=1.02-1.03];  $p < 0.001$ ).

## DISCUSSION

Using the National Hospital Discharge Survey (NHDS) database, we identified six independent risk factors of in-hospital mortality in acute pancreatitis patients: four or more comorbidities, larger hospital size, treatment outside the Midwest region, male sex, older age, and longer hospital length of stay.

Our finding that AP patients with four or more comorbidities are at increased risk of mortality is in agreement with earlier studies [15, 16, 17]. Singh *et al.* found a number of associations between various comorbid conditions and mortality in patients with interstitial pancreatitis [15]. Frey *et al.* reported that an increasing number of chronic comorbidities were associated with higher mortality for acute pancreatitis [16]. These authors also noted that older patients were more likely to die during

**Table 1.** Characteristics evaluated for differences in mortality in acute pancreatitis.

Characteristic	Patients who did not die in the hospital	Patients who died in the hospital	P value
Number of patients	26686	573	
Age (years) (mean ± SD)	64.52 ± 19.11	51.76 ± 19.09	<0.01
Sex			
Male (percent)	13195 (97.8%)	303 (2.2%)	0.1
Female (percent)	13491 (98%)	270 (2.0%)	
Race			
Caucasian (percent)	13292 (97.7%)	318 (2.3%)	0.55
African-Americans (percent)	4875 (27.8%)	109 (2.2%)	
Length of Stay (days) (mean ± SD)	12.02 ± 16.28	6.37 ± 7.96	<0.001
Number of comorbidities			
Three or less	4281 (99.9%)	4 (0.1%)	<0.001
Four or more	22405 (97.5%)	569 (2.5%)	
Region			
Northeast	4934 (97.5%)	129 (2.5%)	0.002
Midwest	7909 (98.4%)	131 (1.6%)	
South	10470 (97.8%)	231 (2.2%)	
West	3373 (97.6%)	82 (2.4%)	
Number of beds in hospital			
6-99	3869 (98.7%)	50 (1.3%)	<0.001
100-199	6266 (97.9%)	134 (2.1%)	
200-299	6346 (98%)	131 (2%)	
300-499	7200 (97.6%)	177 (2.4%)	
>500	3005 (97.4%)	81 (2.6%)	
Hospital Ownership			
Proprietary	2606 (98%)	52 (2%)	0.49
Government	2606 (97.6%)	64 (2.4%)	
Nonprofit or church	21474 (97.9%)	457 (2.1%)	
Principal forms of payment			
Workers compensation	15 (100%)	0 (0%)	<0.001
Medicare	8436 (96.4%)	316 (3.6%)	
Medicaid	4177 (98%)	85 (%)	
Other government	422 (98.4%)	7 (1.6%)	
Blue cross/Blue shield	2576 (98.9%)	29 (1.1%)	
HMO/PPO	4429 (98.6%)	61 (1.4%)	
Other Private Insurance	2747 (98.8%)	33 (1.2%)	
Self-pay	2565 (99%)	27 (1%)	
No charge	204 (99.5%)	1 (0.5%)	
Other	746 (99.2%)	6 (0.8%)	
Not stated	369 (97.9%)	8 (2.1%)	
Type of Admission			
Emergency	18000 (97.9%)	382 (2.1%)	0.19
Urgent	4210 (98.2%)	98.2 (1.8%)	
Elective	1681 (97.2%)	48 (2.8%)	
Not available	2794 (97.7%)	66 (2.3%)	

their hospitalization with AP. An epidemiological study conducted in Japan found that patients older than 70 years with severe comorbidities (>3 comorbid conditions) had twice the risk of in-hospital mortality [17]. Also, Murata *et al.* showed that cardiovascular and renal diseases are factors that increase hospital length of stay and mortality among AP patients [14]. With more comorbid conditions, patients increase their likelihood of developing end organ damage [18]. In our study, patients in the in-hospital mortality group had a longer LOS. However, due to multiple pre-existing comorbidities, those with longer LOS may have had more complications. Others have found that in-hospital mortality was higher in obese patients [19].

We found that in-hospital mortality varied by geographical location. Midwestern states had a lower mortality rate compared to Southern, Western, and Northeastern states. These regional differences amongst hospitals may be related to demographic (e.g., age), clinical (e.g., obesity, number of comorbidities), and socioeconomic (e.g., poverty) factors, population size, and number of medical facilities per capita.

The scoring systems currently utilized to predict mortality in patients with acute pancreatitis are often inconvenient to calculate and do not facilitate efficient triaging. Investigators have used serum markers and radiological findings to predict AP severity and mortality.

**Table 2.** Multivariable analysis of predictors of in-hospital mortality in acute pancreatitis.

Predictor	Adjusted Odds Ratio	95% Confidence Interval	P value
Age	1.03	1.03-1.04	<0.001
Sex-male	1.26	1.05-1.52	0.015
Race			
White	1.28	0.96-1.71	0.09
African-American	1.09	0.87-1.38	0.44
Comorbidities	15.22	4.87-47.54	<0.001
Admission type			
Elective	1.4	0.95-2.07	0.09
Emergency	1.17	0.89-1.55	0.26
Admission source			
Emergency room	0.9	0.7-1.15	0.4
Hospital length of stay	1.03	1.02-1.03	<0.001
Region- Midwest	1.31	1.05-1.64	0.017
Hospital size >100 beds	1.42	1.04-1.95	0.029
Hospital ownership			
Government	1.48	0.97-2.25	0.07
Nonprofit/church	1.2	0.87-1.67	0.27
Principal payment			
Medicare	1.11	0.87-1.43	0.4

To our knowledge, no prior research has used the NHDS database to examine demographic factors, comorbidities, and health care system characteristics that may be related to in-hospital mortality in patients with acute pancreatitis.

While type of hospital ownership (proprietary, government, or owned by a non-profit organization or church) had similar rates of in-patient mortality, smaller hospitals had lower mortality rates. Smaller hospitals may manage primarily milder cases of acute pancreatitis with patients having severe pancreatitis and related complications transferred to large facilities more capable of handling critically ill patients. Future investigations of facility transfer related to pancreatitis may reveal why mortality is lower in smaller hospitals.

Additionally, patients with Medicare as their primary insurance had higher in-hospital mortality compared to other forms of payment. However, the Medicare/mortality relationship found with the univariate analysis was not significant in the multivariable model. As mentioned earlier, unfavorable demographic, clinical, and socioeconomic factors may be more prevalent in Medicare patients than with those having commercial and private insurance coverage.

Mortality among elective and emergency admissions was marginally higher than urgent admissions, but neither hospitalization for elective or emergency reasons was an independent predictor when urgent admission was used as the reference standard. Emergency admissions receive aggressive treatment and higher acuity of care once admitted in comparison to those admitted electively. Without access to individual patient data, we were not able to examine clinical characteristics that might be related to the type of admission.

Our study has several limitations. First, only non-federal hospitals were included in the National Hospital Discharge

Survey database; thus, generalizability to Veterans Affairs medical centers and other institutions should be done with caution. Second, the NHDS database included readmissions for acute pancreatitis; thus, the unit of analysis for the study was events of AP and not independent patients with AP. Finally, the NHDS data did not include lab values, treatment choices, or longitudinal follow-up information; thus, the impact of clinical characteristics could not be examined.

## CONCLUSION

Our analysis of National Hospital Discharge Survey data shows that demographic factors (sex and age), clinical characteristics (comorbidities, length of stay), and features of the health care system (geographic location, hospital size) may be utilized as predictors of in-patient mortality for those diagnosed with acute pancreatitis. Future research may identify other predictors of mortality that would improve the scoring systems now used with acute pancreatitis patients. As more research is conducted, a more accurate, less cumbersome, and easily calculated scoring system may be developed aiding in the reduction of in-hospital mortality in patients with acute pancreatitis.

## Conflict of Interest

We declare that we have no conflict of interests.

## References

1. Vege SS, Yadav D, Chari ST. Pancreatitis. In: GI Epidemiology, 1st ed, Talley NJ, Locke GR, Saito YA (Eds), Blackwell Publishing, Malden, MA 2007.
2. Peery AF, Dellon ES, Lund J, Crockett SD, McGowan CE, Bulsiewicz WJ, et al. Burden of Gastrointestinal Disease in the United States: 2012 Update. *Gastroenterology* 2012; 143: 1179-87.e1-3. [PMID: 22885331]
3. Fagenholz PJ, Castillo CF, Harris NS, Pelletier AJ, Camargo CA Jr. Increasing United States Hospital Admissions for Acute Pancreatitis, 1988-2003. *Ann Epidemiol* 2007; 17:491-7. [PMID: 17448682]

4. Working Group IAP/APA Acute Pancreatitis Guidelines. IAP/ APA Evidence-based Guidelines for the Management of Acute Pancreatitis. *Pancreatology* 2013; 13:e1-15. [PMID: 24054878]
5. Banks PA, Freeman ML; Practice Parameters Committee of the American College of Gastroenterology. Practice guidelines in acute pancreatitis. *Am J Gastroenterol* 2006; 101:2379-400. [PMID: 17032204]
6. Mounzer R, Langmead CJ, Wu BU, Evans AC, Bishehsari F, Muddana V, et al. Comparison of Existing Clinical Scoring Systems to Predict Persistent Organ Failure in Patients With Acute Pancreatitis. *Gastroenterology* 2012; 142:1476-482. [PMID: 22425589]
7. Wang X, Xu Y, Qiao Y, Pang X, Hong L, Fu J, et al. An Evidence-Based Proposal for Predicting Organ Failure in Severe Acute Pancreatitis. *Pancreas* 2013; 42:1255-261. [PMID: 24152952]
8. De Bernardinis M, Violi V, Roncoroni L, Boselli AS, Giunta A, Peracchia A. Discriminant Power and Information Content of Ranson's Prognostic Signs in Acute Pancreatitis: A Meta-analytic Study. *Crit Care Med* 1999; 27:2272-83. [PMID: 10548220]
9. Larvin M. Assessment of clinical severity and prognosis. In: *The Pancreas*, Beger HG, Warshaw AL, Buchler MW, et al (Eds), Blackwell Science, Oxford 1998. p.489.
10. Papachristou GI, Muddana V, Yadav D, O'Connell M, Sanders MK, Slivka A, et al. Comparison of BISAP, Ranson's, APACHE-II, and CTSI Scores in Predicting Organ Failure, Complications, and Mortality in Acute Pancreatitis. *Am J Gastroenterol* 2010; 105:435-41. [PMID: 19861954]
11. Wu BU, Johannes RS, Sun X, Conwell DL, Banks PA. Early Changes in Blood Urea Nitrogen Predict Mortality in Acute Pancreatitis. *Gastroenterology* 2009; 137:129-35. [PMID: 19344722]
12. Mofidi R, Duff MD, Wigmore SJ, Madhavan KK, Garden OJ, Parks RW. Association between Early Systemic Inflammatory Response, Severity of Multiorgan Dysfunction and Death in Acute Pancreatitis. *Br J Surg* 2006; 93:738-44. [PMID: 16671062]
13. Premkumar R, Phillips AR, Petrov MS, Windsor JA. The clinical relevance of obesity in acute pancreatitis: targeted systematic reviews. *Pancreatology* 2015; 15:25-33. [PMID: 25464938]
14. Balthazar EJ, Robinson DL, Megibow AJ, Ranson JH. Acute Pancreatitis: Value of CT in Establishing Prognosis. *Radiology* 1990; 174:331-36. [PMID: 2296641]
15. Singh VK, Bollen TL, Wu BU, Repas K, Maurer R, Yu S, et al. An assessment of the severity of interstitial pancreatitis. *Clin Gastroenterol Hepatol* 2011; 9:1098-1103. [PMID: 21893128]
16. Frey C, Zhou H, Harvey D, White RH. Co-morbidity is a strong predictor of early death and multi-organ system failure among patients with acute pancreatitis. *J Gastrointest Surg* 2007; 11:733-742. [PMID: 17417710]
17. Murata A, Ohtani M, Muramatsu K, Matsuda S. Influence of comorbidity on outcomes of older patients with acute pancreatitis based on a national administrative database. *Hepatobiliary Pancreat Dis Int* 2015; 14:422-8. [PMID: 26256088]
18. Li XY, Wang XB, Liu XF, Li SG. Prevalence and risk factors of organ failure in patients with severe acute pancreatitis. *World J Emerg Med* 2010; 1:201-204. [PMID: 25214968]
19. Murata A, Ohtani M, Muramatsu K, Kobori S, Tomioka S, Matsuda S. Impact of obesity on outcomes of paediatric acute pancreatitis based on a national administrative database. *Pediatr Obes* 2015; 11:174-80. [PMID: 26061540]